

## Medical practice variations

### Appendectomy

Analysis of the distribution and evolution of medical practice in Belgium, in terms of volume and expenditure per insured (analysis and trends by region, province and district), for the year **2021**



NIHDI – Healthcare Service – Directorate for Research, Development and Quality promotion

#### **Appropriate care unit**

Pascal Meeus, Virginie Dalcq, Delphine Beauport, Katrien Declercq, Lucien Hoekx, Kathleen Sierens, Benjamin Swine

Contact: [appropriatecare@riziv-inami.fgov.be](mailto:appropriatecare@riziv-inami.fgov.be)

**Date of report: 8 June 2022**

## CONTENTS

<b>CONTENTS .....</b>	<b>2</b>
<b>1. INTRODUCTION .....</b>	<b>3</b>
<b>2. SPECIFIC METHOD OF ANALYSIS .....</b>	<b>4</b>
A. NIHDI NOMENCLATURE CODES SELECTED FOR ANALYSIS.....	4
B. PAST HISTORY OF NOMENCLATURE CODES .....	5
C. SOURCE OF DATA AND ANALYSIS PERIOD.....	6
D. SPECIFIC SELECTION CRITERIA .....	7
E. STANDARDISATION.....	7
<b>3. RESULTS .....</b>	<b>8</b>
A. NATIONAL STANDARDISED RATE OF USE.....	8
B. BREAKDOWN OF NOMENCLATURE CODES PROVIDED, BY VOLUME.....	9
C. SPECIALISATION OF HEALTHCARE PROVIDERS .....	10
D. SPECIALISATION OF PRESCRIBERS.....	11
E. STANDARDISED RATE OF USE BY SEX AND AGE GROUP .....	12
F. STANDARDISED RATE OF USE: HOSPITAL AND OUTPATIENT CARE .....	16
G. STANDARDISED RATE OF USE BY REIMBURSEMENT SCHEME .....	18
H. TRENDS IN STANDARDISED RATES OF USE .....	20
I. GEOGRAPHICAL VARIATIONS IN STANDARDISED RATES OF USE .....	24
J. STANDARDISED HEALTHCARE EXPENDITURE BORNE BY THE INSURANCE .....	29
<b>4. KEY DATA SUMMARY .....</b>	<b>32</b>
<b>5. APPENDICES .....</b>	<b>33</b>
A. ANALYSIS OF VARIANCE (ANOVA), EXCEPT BRUSSELS .....	33
B. FREQUENCY OF PRACTICE OCCURRENCES.....	34
C. PATIENT CARE SETTINGS.....	36
D. CODING VARIATIONS AND PRACTICE ALTERNATIVES.....	38

### 1. INTRODUCTION

The Appropriate Care Unit was set up within the NIHDI's Directorate for Research, Development and Quality under NIHDI's Administration Contract for 2016-2018<sup>1</sup>. Article 35 of this contract refers to 'the setting up of an Appropriate Care Unit, aiming specifically to promote an integrated approach to the rational use of resources'. The Appropriate Care Unit has been up and running since the second quarter of 2017.

The tasks of the Unit were set out formally in the '2016-2017 Healthcare monitoring Action plan', published by NIHDI on 18 July 2016<sup>2</sup>. This plan lists around thirty measures designed to make healthcare provision more efficient, by encouraging appropriate practice and tackling unnecessary or inappropriate care.

The plan states that one of the tasks of the Appropriate Care Unit is to analyse the 'appropriateness of care', in order to identify unexplained variations in consumption patterns, identified after standardisation. Such variations can potentially point to non-optimal use of resources.

'Variations in medical practice' documents report on the analyses carried out in this framework. Each report focuses on a particular topic.

In this document, we present the figures and graphs relating to analyses<sup>3</sup> of practice in the area of Appendectomy, and give the explanations necessary to understand these.

We have deliberately chosen not to attempt to interpret the figures, preferring to present the results to experts who are in a better position to do so. This document has nevertheless been made available to the public in order to provide objective, open input to discussions on this issue.

---

<sup>1</sup> (Institut national d'assurance maladie-invalidité, 2016)

<sup>2</sup> (Institut national d'assurance maladie-invalidité, 2016)

<sup>3</sup> Readers interested in the methodology used in these quantitative analyses should consult the document entitled 'Variations in practice – Methodology'.

## 2. SPECIFIC METHOD OF ANALYSIS

### A. NIHDI nomenclature codes selected for analysis

The NIHDI nomenclature codes selected for the analysis are listed below:

Outpatient	Inpatient	Rates	Expenses	Label	Creation	Deletion	Group N	Value
156310	156321	no	yes	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 243154-243165 de la nomenclature, par voie endoscopique	01-07-2014		N80	.
156332	156343	no	yes	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 244871-244882 de la nomenclature, par voie endoscopique	01-07-2014		N80	.
156354	156365	no	yes	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 244871-244882 de la nomenclature, en chirurgie ouverte	01-07-2014		N80	.
243154	243165	yes	yes	Appendicectomie à froid ou à chaud, avec ou sans perforation	01-04-1985		N23	N200
243353	243364	yes	yes	Appendicectomie à froid ou à chaud, avec ou sans perforation	01-10-1995	31-12-2011	N23	.
244871	244882	yes	yes	Appendicectomie nécessitant la résection du fond caecal	01-05-2007		N23	N400
731032	731043	no	yes	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 243154-243165, par voie endoscopique	01-03-2004	30-06-2014	N80	.
736750	736761	no	yes	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 244871-244882, par voie endoscopique	01-04-2011	30-06-2014	N80	.
736772	736783	no	yes	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 244871-244882, en chirurgie ouverte	01-04-2011	30-06-2014	N80	.

This table shows the NIHDI nomenclature codes selected for this analysis, stating whether or not they were included in the analyses of services and expenditure, and giving, for each one, a description, dates of creation and deletion, where appropriate, their N group (in the NIHDI nomenclature) and their value.

## B. Past history of nomenclature codes

Outpatient	Inpatient	Date	Label
731032	731043	01-03-2004	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 243353 - 243364, par voie endoscopique
731032	731043	01-01-2012	Ensemble du matériel de consommation et du matériel implantable utilisé lors de la prestation 243154-243165, par voie endoscopique



This table displays the historic evolution of the definitions of the NIHDI-nomenclature codes taken into account for this analysis, if modifications were implemented during the period 2011-2021.

### C. Source of data and analysis period

The data used in the analyses have been taken from the following databases:

<b>Document N</b>	for the utilisation rate and amount of expenses of insured persons (who meet the selection criteria) whose age, sex, preferential regime and residence are known 2011-2021
<b>Document P</b>	for the utilisation rate and amount of expenses of insured persons (who meet the selection criteria) by type of medical specialities in 2021 Semester 1 (data extrapolated to the full year)
<b>Document P, SHA, ADH</b>	for the practice occurrences and analysis of patient care settings in 2019
-	-

<b>Analysis period</b>	2011-2021
------------------------	-----------



'N Documents' are monthly data sent by the sickness funds to NIHDl, within three months. These data show the number of services provided, dates and the fees involved. Every six months, these data are compiled by the insurers, which also add data on patients: age, gender, social category and district of residence. N Documents, however, cannot be used to analyse the combinations of services received by individual patients.

'P Documents' are six-monthly data sent by the sickness funds to NIHDl, within four months. These data show the services provided, the service-provider, the prescriber, the place of provision of service, and the hospital where patients were treated. P Documents can be used to monitor medical consumption and pricing, but not (yet) to analyse services per patient.

'Documents SHA, ADH' are sent annually and within six months by the insurer-organisations to the NIHDl. They include all the services provided respectively in day admission and standard hospitalisation, in general hospitals per hospital stay.

#### D. Specific selection criteria

Several filters may have been applied to the data, so that only one section of the population is considered in the analyses. If so, the filters used are shown in the table below:

FILTERS APPLIED TO DATA	
Sex	women and men
Age	all
-	-

#### E. Standardisation

The data are standardised before analysis per year, based on age, sex and preferential regime per arrondissement, province and region (standardization based on population in 2021).



**Standardisation** renders populations comparable in relation to one or several criteria. If a difference is observed between these populations, we can therefore assume that it is not due to the criteria covered by the standardisation process.


### 3. RESULTS

#### A. National standardised rate of use

	TOTAL
Average number of interventions per year	14.892
Standardised rate of use per 100 000 insured persons	130



## B. Breakdown of nomenclature codes provided, by volume



See page 4 for details about the NIHD nomenclature codes selected for analysis.

Note : The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis.

## C. Specialisation of healthcare providers

Specialisation of the provider*	Total providers	Concerned providers	% Providers	Median of H.C. services	Q3 of H.C. services	% Total H.C. services
General surgery	1307	487	37%	26	42	99,39%
Other specialities	4807	33	1%	2	3	0,61%
Total	6114	520	9%	24	40	100,00%



This table shows the following non-standardised data, by medical specialities (figures for the year 2021):

- The number of service-providers per specialisation who have recorded at least one service (the figures are exceptionally extrapolated from a single semester if an \* is indicated in the header, otherwise the full year is taken into account);
- The number of service-providers recording services under the nomenclature codes selected for this analysis;
- The service-providers for these codes as a percentage of the total number of service-providers recording provision of at least one service;
- The median number and third quartile of services per service-provider (recording provision under these codes);
- The service percentage, i.e. the number of services recorded for this specialisation as a percentage of total services provided.

## D. Specialisation of prescribers

Specialisation of the prescriber*	Total prescribers	Concerned prescribers	% Prescribers	Median of prescriptions	Q3 of prescriptions	% Prescriptions
Not applicable	0	0	0%	0	0	100,00%
Total	0	0	0%	0	0	100,00%



This table shows, in order, the following non-standardised data per specialities (figures for the year 2021):


- The number of prescribers who have prescribed at least one service (the figures are exceptionally extrapolated from a single semester if an \* is indicated in the header, otherwise the full year is taken into account);
- The number of prescribers prescribing the nomenclature codes selected for this analysis;
- The prescribers prescribing these codes as a percentage of the number of prescribers prescribing at least one service;
- The median number and third quartile of services per prescriber (prescribing these codes);
- The percentage of services prescribed, i.e. the number of prescriptions issued for this specialisation as a percentage of total services prescribed.

## E. Standardised rate of use by sex and age group

	TOTAL
<i>Average number of interventions per year</i>	<b>14.892</b>
Median age (years)	<b>30</b>
Mean age (years)	<b>34,02</b>
Max/Min Ratio of the median age (by district)	<b>1,86</b>
Percentage of women	<b>47,29%</b>

## Max/Min Ratio:


The max/min ratio measures the dispersion of values. It is calculated as the ratio of the maximum value found for the variable, in all districts, to the minimum value. If this minimum value is equal to zero, the max/min ratio cannot be calculated, and is reported as 'NA' ('not applicable').




This figure is made up of bar charts for each sex. The **coefficient of variation**, shown by the red line, measures the relative dispersion of the standardised rates of use observed for each district, by age group and sex (standard deviation divided by the mean). This line is shown in bold for age groups where the coefficient of variation can be validly interpreted (i.e. for age groups in which there are sufficient insured persons per district to allow for a proper comparison).

The left-hand vertical axis of the graph represents the standardised rate of use, and the right-hand axis the coefficient of variation. The horizontal axis shows the age groups. The horizontal dotted lines show the total values of the standardised rates of use (in blue) and of the coefficient of variation (in red).

Standardised rate of use per 100 000 insured persons, and coefficient of variation for the districts, by age group and sex, for the year 2021



Comparison of the standardised rates of use by sex (per 100 000) in 2021




This histogram shows standardised rates of use by province and by sex. The grey bars show the rates for men, while the green bars show the rates for women, for each province. The grey and green broken lines show the total standardised rates of use, again grey for men, green for women.

Standardised rate of use per 100 000 insured persons, by sex and by province for the year 2021

## F. Standardised rate of use: hospital and outpatient care


	TOTAL
<i>Average number of interventions per year</i>	<b>14.892</b>
Percentage of out-patient care	<b>2,91%</b>
Max/min ratio of out-patient care percentage (by district)	NA (Min=0)



Percentage of outpatient care, total and by region

This graph shows the percentage of outpatient services (including hospital day admissions), i.e. the number of outpatient services provided as a percentage of total services (outpatient and hospital stays). Besides the bar per region, there is a bar for the entire Belgian population. A dotted line also shows this overall ratio.






Change over time in the percentage of outpatient care, by province

N.B.:

- The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis
- A complementary document to this chapter, about the handling of patients per health care sector, is [enclosed in this report](#) (cf. p.36)

## G. Standardised rate of use by reimbursement scheme

	TOTAL
<i>Average number of interventions per year</i>	<b>14.892</b>
Percentage provided under the preferential reimbursement scheme	<b>16,13%</b>
Standardised rate of use with preferential reimbursement scheme (per 100 000)	<b>115</b>
Standardised rate of use without preferential reimbursement scheme (per 100 000)	<b>133</b>
Ratio Preferential scheme /General scheme	<b>0,86</b>




Standardised rate of use by reimbursement scheme and by region

This graph shows the standardised rates of use with (in red) and without (in grey) the preferential reimbursement scheme, by region and in total. The red and grey dotted lines show the overall standardised rates of use, with and without the preferential reimbursement scheme, respectively.

## H. Trends in standardised rates of use

	TOTAL
<i>Average number of interventions per year</i>	<b>14.892</b>
<b>Trend (2011-2021)</b>	<b>0,25%</b>
Trend (2011-2019)	-0,59%
Trend (2019-2021)	3,69%


These trends correspond to the average annual growth rate.



This graph shows a coloured curve for each region and a black curve for the entire Belgian population. The x-axis shows the years, and the y-axis shows the standardised rate of use per 100 000 insured persons.

Note : The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis.

Trends in the standardised rate of use per 100 000 insured persons, by region



Trends in the standardised rate of use per 100 000 insured persons, by district

This graph shows a colored line for each district and a black line for the entire Belgian population. The x-axis shows the years, and the y-axis shows the standardised rate of use per 100 000 insured persons.

To better highlight changes over time, the rates shown are **rolling averages** of the rates for the three years preceding the year in question (including the year itself).

The graph only shows the five districts with the highest average rates and the five districts with the lowest average rates over the last 3 years studied.

Note : The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis.

		Rate of use	Annual increase			Structural break
		2021 (per 10 <sup>5</sup> insured)	2011-2021	2011-2019	2019-2021	
Provinces	West Flanders	146,97	-0,15%	-0,44%	1,03%	NA
	East Flanders	138,23	0,16%	-1,05%	5,14%	NA
	Antwerp	126,42	0,71%	0,23%	2,66%	NA
	Limburg	135,71	0,79%	-0,19%	4,83%	NA
	Flemish Brabant	144,57	1,50%	-0,44%	9,61%	NA
	Brussels	109,96	-0,50%	-1,63%	4,13%	NA
	Walloon Brabant	127,47	-1,40%	-0,80%	-3,75%	NA
	Hainaut	115,74	-0,04%	-0,15%	0,44%	NA
	Liège	128,43	0,57%	-0,61%	5,41%	NA
	Namur	121,84	-0,42%	-1,31%	3,26%	NA
	Luxembourg	127,09	-0,60%	-1,53%	3,22%	NA
Regions	Flanders	137,09	0,55%	-0,36%	4,30%	NA
	Brussels	109,96	-0,50%	-1,63%	4,13%	NA
	Wallonia	122,60	-0,12%	-0,61%	1,89%	NA
TOTAL		129,89	0,25%	-0,59%	3,69%	NA


Trends in the rates of use, by province and region

This table reports the standardised **rates of use** for the last year analysed (2021), as well as the average **rates of increase**, by province, by region and in total, for the entire period (2011-2021), for the last three years (2019-2021) and for the period preceding the last three years (2011-2019)

In order to find out whether the trend in the last three years differs from that in the years before, a linear mixed model was fitted in two steps. In the first step a change in trend on the national level is tested. If this test is significant, in a second step, the model tests whether the difference in trend is significant for each province, region and at the national level. The data of 2020 are excluded from the models.

The significance of the test for a change in trend is reported in the Structural break column : \* P-value ≤ 0.05 / \*\* P-value ≤ 0.01 / \*\*\* P-value ≤ 0.001 and NS for a non-significant result.

'NA' is shown where the nomenclature codes selected for the analysis have not been used for the entire three-year period or when the statistical tests cannot be evaluated.



Trend break assessment model by province – Regression lines

Regression lines per province showing a possibly different slope for the last three years (2019-2021) compared to the years before (2011-2019).

Data of 2020 was excluded from this analysis, but is indicated on the graph for information.


## I. Geographical variations in standardised rates of use

	TOTAL
<i>Average number of interventions per year</i>	<b>14.892</b>
<b>Coefficient of Variation (2021)</b>	<b>15,48</b>
<b>Max/Min Ratio* of the standardised rates of use (by region)</b>	<b>1,25</b>
<b>Max/Min Ratio* of the standardised rates of use (by district)</b>	<b>2,6</b>

<b>Coefficient of Variation (2019-2021)</b>	<b>14,41</b>
<b>Coefficient of Variation (2011-2013)</b>	<b>11,73</b>
<i>Statistically significant difference? (<math>p \leq 0.05</math>)</i>	<i>No</i>

\* An 'NA' result indicates a ratio which cannot be calculated, i.e. the minimum value = zero (cf. E. Standardised rate of use by sex and age group)






'Dot plot' showing standardised rates of use by district, by sex

A **dot plot** is a distribution chart, which is useful for highlighting groups in the data, gaps in the distribution and outliers. Here, each dot represents the rate of use of a district, for its entire population or broken down by sex.

The rates are rounded to the nearest unit, ten, hundred, etc., depending on the value of the maximum rate, in order to better group the values.

The graph also shows a box with the 25<sup>th</sup>, 50<sup>th</sup> and 75<sup>th</sup> percentiles of the non-rounded standardised rates of use for all patients. The bottom line of the box represents the 25<sup>th</sup> percentile, while the upper line represents the 75<sup>th</sup> percentile. The line inside the box represents the 50<sup>th</sup> percentile.




Map showing distribution of standardised rates of use, by district

On this map of Belgium, thin lines show the boundaries of the districts, while thick lines show the provincial borders. The districts are coloured using a colour scale based on the level of rate of use in the district compared to the Belgian national rate (overall rate). This ratio is expressed as a percentage: e.g. 0% if the district rate is equal to the overall rate, 20% if the rate is 20% above the overall rate, and -20% if the rate is 20% below the overall rate. The percentages are calculated using the standardised rates of the last year analysed, and are displayed in bands of 20%. The following colour coding applies:

Colour	Category
Dark red	More than 50%
Red	Between 30% and 50%
Orange	Between 10% and 30%
Yellow	Between -10% and 10%
Light green	Between -30% and -10%
Green	Between -50% and -30%
Dark green	Less than -50%
White	Not used


N.B.: The interpretation of this map is to be done in parallel with [the graph in funnel plot](#) (p.28)



Map showing distribution of standardised expenditure, by district

On this map of Belgium, thin lines show the boundaries of the districts, while thick lines show the provincial borders. The districts are coloured using a colour scale based on the level of expenditure in the district compared to Belgian national (overall) expenditure. This ratio is expressed as a percentage: e.g. 0% if expenditure in the district is equal to the overall expenditure, 20% if it is 20% higher, and -20% if it is 20% lower. The percentages are calculated using the standardised expenditure of the last year analysed and are displayed in bands of 20%. The following colour coding applies:

Colour	Category
Dark red	More than 50%
Red	Between 30% and 50%
Orange	Between 10% and 30%
Yellow	Between - 10% and 10%
Light green	Between -30% and -10%
Green	Between -50% and - 30%
Dark green	Less than -50%
White	No expenditure



In this graph, the standardised rate of use in a district is positioned versus the size of its population. Besides the dots representing the districts, 95% and 99.7% **confidence intervals** are also shown on the graph. These are dependent of the size of the districts. The thicker horizontal line shows the national standardised rate of use. The outlier districts are identified as those districts that fall outside the 99.7% confidence intervals, the zone between the 95% and 99.7% confidence intervals being considered as "warning zone".

N.B.: The interpretation of this graph is to be done in parallel with the [map of the distribution of rates of use](#) (p.26)

**'Funnel plot' showing the standardised rates of use by district, by the number of insured persons**

## J. Standardised healthcare expenditure borne by the insurance

	TOTAL
<i>Average number of interventions per year</i>	<b>14.892</b>
<b>Average annual expenditure (€)</b>	<b>8.332.102</b>
<b>Average cost per intervention (€)</b>	<b>559,5</b>
<b>Average annual expenditure per insured (€)</b>	<b>0,73</b>
<b>Max/Min Ratio* of expenditure per insured (by region)</b>	<b>1,32</b>
<b>Max/Min Ratio* of expenditure per insured (by district)</b>	<b>3,41</b>

\* An 'NA' result indicates a ratio which cannot be calculated, i.e. the minimum value = zero (cf. E. Standardised rate of use by sex and age group)

		Standardised expenditure (per insured)
Provinces	West Flanders	0,79 €
	East Flanders	0,81 €
	Antwerp	0,72 €
	Limburg	0,72 €
	Flemish Brabant	0,87 €
	Brussels	0,59 €
	Walloon Brabant	0,65 €
	Hainaut	0,62 €
	Liège	0,74 €
	Namur	0,62 €
	Luxembourg	0,79 €
Regions	Flanders	0,78 €
	Brussels	0,59 €
	Wallonia	0,67 €
TOTAL		0,73 €

Regional and provincial distribution of standardised expenditure (2021)

Nomenclature	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	Average annual growth rate
243154-243165	232,63	242,08	245,67	249,38	249,66	249,78	251,86	255,99	256,33	258,45	260,34	1,13%
243353-243364	239,50	239,75	228,65	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	NA
244871-244882	457,26	465,88	473,95	482,11	480,20	480,94	484,65	491,32	492,64	499,46	503,16	0,96%

Change over time in expenditure, by service and by nomenclature code

## 4. KEY DATA SUMMARY

		TOTAL	
PROVIDERS & PRESCRIBERS			
Main healthcare providers:		General surgery	99,39%
Main prescribers:		Not applicable	-
RATE OF USE			
Average number of interventions (per year)			14.892
Standardised rate of use (per 100 000 insured persons)			129,89
≥ 2 occurrences per patient (2019) <sup>4</sup>			0,6%
Percentage of outpatient care			2,91%
POPULATION			
Median age			30 years
Max/min ratio <sup>5</sup> of the median age (by district)			1,86
Percentage of women			47,29%
Ratio Preferential rate/General rate			0,86
TRENDS			
Trend (2011-2021)			0,25%
Trend <sup>6</sup> (2011-2019)			-0,59%
Trend <sup>6</sup> (2019-2021)			3,69%
GEOGRAPHICAL VARIATIONS			
Coefficient of variation <sup>6</sup> (2011-2013)			11,73
Coefficient of variation <sup>6</sup> (2019-2021)			14,41
Max/min <sup>5</sup> Ratio of number of interventions <sup>6</sup> (per 100 000 insured persons, by region)			1,25
Max/min Ratio <sup>5</sup> of number of interventions (per 100 000 insured persons, by district)			2,6
DIRECT EXPENDITURE			
Average annual expenditure			8.332.102 €
Average annual expenditure per insured			0,73 €
Max/Min Ratio <sup>5</sup> of expenditure per insured (by region)			1,32
Max/Min Ratio <sup>5</sup> of expenditure per insured (by district)			3,41
Average cost of interventions			559,5 €
CODING VARIATIONS & PRACTICE ALTERNATIVES <sup>4</sup>			
Variations in practice coding <sup>6</sup> (by province)			Yes
Variations in the choice of practice alternatives <sup>6</sup> (by province)			Yes

<sup>4</sup> More detailed results are shown in a document enclosed to this report.

<sup>5</sup> An 'NA' result indicates a ratio, which cannot be calculated, i.e. the minimum value equals zero.

<sup>6</sup> If the result(s) show(s) a significant difference, the level of statistical significance is symbolized by one to three asterisks (increasingly significant). Otherwise, NS is displayed (not significant). 'NA' indicates the test is not applicable.



## 5. APPENDICES

### A. Analysis of variance (ANOVA), except Brussels

Statistical significance of the differences observed in 2021		
<i>By region?</i>	Yes	*
<i>By sex?</i>	Yes	*
<i>By reimbursement scheme?</i>	Yes	*
<i>By sex and per region?</i>	No	NS
<i>By reimbursement scheme and per region?</i>	No	NS
<i>By sex and per reimbursement scheme?</i>	No	NS
<i>By sex and reimbursement scheme and per region?</i>	No	NS


In order to be able to assess the significance of the observed differences, a linear mixed **ANOVA** model was fitted to the data of all districts of the Walloon and Flemish regions, after standardising for age. The model has region, sex and reimbursement scheme as fixed effects and also contains all two-way and three-way interactions between these effects.

In order to interpret the model correctly, first the three-way interaction should be evaluated, followed by the two-way interactions and finally by the main effects. If the three-way interaction is significant, the interpretation of the model should be done at this level only and the two-way interactions and main effects should not be interpreted. If the three-way interaction is not significant, the two-way interactions are evaluated. Every main effects that appears in a significant interaction should be interpreted at the level of the interaction and not at the level of that main effect. Main effects can only be interpreted directly if they don't appear in a significant interaction.

The **asterisks** represent the level of statistical significance of the tests: \* P-value  $\leq 0,05$  / \*\* P-value  $\leq 0,01$  / \*\*\* P-value  $\leq 0,001$  or NS for a non-significant result.

## B. Frequency of practice occurrences

Frequency	Per year	Per day
2 occurrences	0,5%	0,4%
≥ 3 occurrences	0,1%	0,1%
≥ 2 occurrences	0,6%	0,5%




Distribution of practice recurrences per year (2019)

Some practices may be billed several times for the same patient in the same year or even on the same day. This may be due to a **repetition of the practice**, but also to an anatomical effect, which may lead, depending on the organ concerned, to performing the same practice **bilaterally**, which may therefore cause a double occurrence on the same day.

In order to interpret the results per day validly, it is useful to note that the same patient may be counted several times if, for example, he or she has received two identical services simultaneously, twice a year.


These frequency analyses of occurrences are carried out over the year **2019** using the following databases: Documents P, ADH and SHA.

Values « **n.a.** » are indicated if the data were not available at the time of this report.



Frequency of practice occurrences by province and variation vs national value (2019)

## C. Patient care settings



■ Inpatient
 ■ One-day
 ■ Outpatient (clinic)


Distribution of patient care settings in 2019

Care Settings	
Outpatient (private)	0,0%
Outpatient (polyclinic)	0,3%
(Day) Hospital	2,3%
Hospital (stay)	97,3%

In addition to the chapter on [standardised inpatient and outpatient use rates](#) (see p.16), the analysis of patient care settings can be refined by identifying the outpatient (private and polyclinic) and inpatient (day or standard hospitalisation) sub-sectors.

These analyses are carried out over the year **2019** using the following databases: Documents P, ADH and SHA.


Values « **n.a.** » are indicated if the data were not available at the time of this report.



Distribution of patient care settings by province (2019)

## D. Coding variations and practice alternatives

### → Variations in coding:




Outpatient	Inpatient	Label
243154	243165	Appendicectomie à froid ou à chaud, avec ou sans perforation
243353	243364	Appendicectomie à froid ou à chaud, avec ou sans perforation
244871	244882	Appendicectomie nécessitant la résection du fond caecal

Significance	By region	By province
Use of Nomenclature codes <sup>7</sup>	***	***

<sup>7</sup> The calculation of significance is carried out here by comparing the geographical differences in the use of the different nomenclature codes to code the practice.

The **asterisks** represent the level of statistical significance of Chi-square test: \* P-value ≤ 0,05 / \*\* P-value ≤ 0,01 / \*\*\* P-value ≤ 0,001. **NS** and **NA** respectively indicate that the variations are not significant or not applicable.

## → Variations in practice alternatives:



Breakdown of choice for practice alternatives


Combined codes	Groupings
<b>243154-243165</b>	1_w/o_caecum
<b>244871-244882</b>	2_w/_caecum

Significance	By region	By province
Choice of Practice alternatives	***	***



According to the nature of the practice and the analytical tools available, it may be possible to identify and define alternatives for carrying out the practice. In this case, the nomenclature codes defined for the analysis of the practice are grouped together with the aim of analysing whether or not the choices of these alternatives are homogeneous across the territory. The calculation of significance displayed in the table is carried out by comparing these groups of codes with each other.

## → Variation in the use of Global Payment with Standardisation (GPS) :



Pseudocodes	Label
-799002	Groupe des patients ayant bénéficié d'une appendectomie, en hospitalisation classique, de sévérité 1.

Breakdown by volume of pseudocodes of GPS

Significance	By region	By province
Use of Global Payment with Standardisation <sup>8</sup>	NS	***

<sup>8</sup> The calculation of significance is carried out here by comparing the use of Global Payment with Standardisation as a whole compared to the non-use of these packages.