

Medical practice variations

Extraction (Adjusted)

Analysis of the distribution and evolution of medical practice in Belgium, in terms of volume and expenditure per insured
(analysis and trends by region, province and district), for the year **2022**



NIHDI – Healthcare Service – Directorate for Research, Development and Quality promotion

Appropriate care unit

Pascal Meeus, Virginie Dalcq, Delphine Beauport, Katrien Declercq, Benjamin Swine

Contact: appropriatecare@riziv-inami.fgov.be

Date of report: 9 August 2023

CONTENTS

CONTENTS	2
1. INTRODUCTION	3
2. SPECIFIC METHOD OF ANALYSIS	4
A. NIHDI NOMENCLATURE CODES SELECTED FOR ANALYSIS.....	4
B. PAST HISTORY OF NOMENCLATURE CODES	5
C. SOURCE OF DATA AND ANALYSIS PERIOD.....	6
D. SPECIFIC SELECTION CRITERIA	7
E. STANDARDISATION.....	7
3. RESULTS	8
A. NATIONAL STANDARDISED RATE OF USE.....	8
B. BREAKDOWN OF NOMENCLATURE CODES PROVIDED, BY VOLUME	9
C. SPECIALISATION OF HEALTHCARE PROVIDERS	10
D. SPECIALISATION OF PRESCRIBERS.....	11
E. STANDARDISED RATE OF USE BY SEX AND AGE GROUP	12
F. STANDARDISED RATE OF USE: HOSPITAL AND OUTPATIENT CARE	16
G. STANDARDISED RATE OF USE BY REIMBURSEMENT SCHEME	18
H. TRENDS IN STANDARDISED RATES OF USE	20
I. GEOGRAPHICAL VARIATIONS IN STANDARDISED RATES OF USE	24
J. STANDARDISED HEALTHCARE EXPENDITURE BORNE BY THE INSURANCE	29
4. KEY DATA SUMMARY	32
5. APPENDICES	33
A. ANALYSIS OF VARIANCE (ANOVA), EXCEPT BRUSSELS	33
B. FREQUENCY OF PRACTICE OCCURRENCES.....	34
C. PATIENT CARE SETTINGS.....	36
D. CODING VARIATIONS AND PRACTICE ALTERNATIVES.....	38

1. INTRODUCTION

The Appropriate Care Unit was set up within the NIHDI's Directorate for Research, Development and Quality under NIHDI's Administration Contract for 2016-2018¹. Article 35 of this contract refers to 'the setting up of an Appropriate Care Unit, aiming specifically to promote an integrated approach to the rational use of resources'. The Appropriate Care Unit has been up and running since the second quarter of 2017.

The tasks of the Unit were set out formally in the '2016-2017 Healthcare monitoring Action plan', published by NIHDI on 18 July 2016². This plan lists around thirty measures designed to make healthcare provision more efficient, by encouraging appropriate practice and tackling unnecessary or inappropriate care.

The plan states that one of the tasks of the Appropriate Care Unit is to analyse the 'appropriateness of care', in order to identify unexplained variations in consumption patterns, identified after standardisation. Such variations can potentially point to non-optimal use of resources.

'Variations in medical practice' documents report on the analyses carried out in this framework. Each report focuses on a particular topic.

In this document, we present the figures and graphs relating to analyses³ of practice in the area of Extraction (Adjusted), and give the explanations necessary to understand these.

We have deliberately chosen not to attempt to interpret the figures, preferring to present the results to experts who are in a better position to do so. This document has nevertheless been made available to the public in order to provide objective, open input to discussions on this issue.

¹ (Institut national d'assurance maladie-invalidité, 2016)

² (Institut national d'assurance maladie-invalidité, 2016)

³ Readers interested in the methodology used in these quantitative analyses should consult the document entitled 'Variations in practice – Methodology'.

2. SPECIFIC METHOD OF ANALYSIS

A. NIHDI nomenclature codes selected for analysis

The NIHDI nomenclature codes selected for the analysis are listed below:

Outpatient	Inpatient	Rates	Expenses	Label	Creation	Deletion	Group N	Value
301151	301162	yes	yes	* Extraction d'une dent, chez un patient à risque hémorragique répondant aux conditions de l'article 6 § bis de la nomenclature, à partir du 5 ^e anniversaire pour autant qu'une suture ait été effectuée	01-01-2018	30-06-2022	N04	.
301173	301184	yes	yes	* Extraction d'une dent chez un patient à risque hémorragique répondant aux conditions de l'article 6 § bis de la nomenclature, à partir du 18 ^e anniversaire jusqu'au 50 ^e anniversaire, pour autant qu'une suture ait été effectuée	01-01-2018	30-06-2022	N04	.
301195	301206	yes	yes	+ Extraction chirurgicale d'une dent environnant et suture des lambeaux muqueux incisés, avec résection de l'os environnant et suture des lambeaux muqueux incisés, à partir du 50 ^e anniversaire	01-01-2018		N04	L63
301210	301221	yes	yes	+ Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, à partir du 50 ^e anniversaire jusqu'au 50 ^e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6 § bis	01-01-2018		N04	L63
304850	304861	yes	yes	+ Extraction d'une dent, y compris la suture éventuelle, à partir du 50 ^e anniversaire	01-06-2007	30-06-2022	N04	.
304872	304883	yes	yes	+ Extraction d'une dent, y compris la suture éventuelle, à partir du 18 ^e anniversaire jusqu'au 50 ^e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6 § bis	01-06-2007		N04	L15
304894	304905	yes	yes	+ Extraction d'une dent, à partir du 50 ^e anniversaire, par dent supplémentaire	01-06-2007	30-06-2022	N04	.
304916	304920	yes	yes	+ Extraction d'une dent, à partir du 50 ^e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6 § bis	01-06-2007		N04	L15
304975	304986	yes	yes	+ Extraction d'une dent, à partir du 50 ^e anniversaire	01-07-2022		N04	L21
304990	305001	yes	yes	+ Extraction d'une dent, à partir du 18 ^e anniversaire jusqu'au 50 ^e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § bis	01-07-2022		N04	L21
305130	305141	no	yes	Honoraires complémentaires pour suture de plaine pendant une séance d'extraction(s) dentaire(s) ou ablation (section et extraction) de racine(s), à l'exception des prestations où la suture est comprise, à partir du 18 ^e anniversaire	01-07-2022		N04	L12
305152	305163	no	yes	Honoraires complémentaires pour suture de plaine pendant une séance d'extraction(s) ou ablation (section et extraction) de racine(s), à l'exception des prestations où la suture est comprise, par dent supplémentaire dans le même quadrant et durant la même séance, à partir du 18 ^e anniversaire	01-07-2022		N04	L8
312410	312421	yes	yes	Opération pour l'ablation d'une dent incluse par résection osseuse périconvolute et, le cas échéant, lorsqu'une section dentaire est effectuée	01-02-2004		N16	K120
312422	312443	yes	yes	Ostéotomie au-dessus et au-dessous d'un gencive dentaire avec section dentaire éventuelle et enlèvement du gencive dentaire	01-02-2004		N16	K120
317214	317225	yes	yes	+ Extraction sous anesthésie générale, de 8 dents ou moins, y compris l'ablation de la carie et les sutures éventuelles	01-04-1985		N16	K125
317236	317240	yes	yes	+ Extraction, sous anesthésie générale, de moins de 8 dents, y compris l'avoléotomie et les sutures éventuelles	01-04-1985		N16	K75
317251	317262	yes	yes	Extraction en milieu hospitalier de minimum 8 dents, y compris l'avoléotomie et les sutures éventuelles, et ce sous monitoring des paramètres vitaux	01-02-2004		N16	K125
317376	317380	yes	yes	Extraction en milieu hospitalier de moins de 8 dents, y compris l'avoléotomie et les sutures éventuelles, et ce sous monitoring des paramètres vitaux	01-02-2004		N16	K75
317391	317402	yes	yes	+ Extraction d'une canine latérale, d'une molaire latérale ou d'une dent définitive, chez un patient à risque hémorragique répondant aux conditions de l'article 6 § bis de la nomenclature, jusqu'au 18 ^e anniversaire, pour autant qu'une suture ait été effectuée	01-02-2004		N16	K75
371151	371162	yes	yes	+ Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, jusqu'au 18 ^e anniversaire	01-01-2018	30-06-2022	N04	.
371195	371206	yes	yes	+ Extraction d'une canine latérale, d'une molaire latérale ou d'une dent définitive, y compris la suture éventuelle, jusqu'au 18 ^e anniversaire	01-01-2018		N04	L63
374850	374861	yes	yes	+ Extraction d'une canine latérale, d'une molaire latérale ou d'une dent définitive, y compris la suture éventuelle, jusqu'au 18 ^e anniversaire	01-06-2007	30-06-2022	N04	.
374872	374883	yes	yes	+ Extraction d'une canine latérale, d'une molaire latérale ou d'une dent définitive, jusqu'au 18 ^e anniversaire, par dent supplémentaire dans le même quadrant et durant la même séance	01-06-2007		N04	L15
374975	374986	yes	yes	Honoraires complémentaires pour suture de plaine pendant une séance d'extraction(s) dentaire(s) ou ablation (section et extraction) de racine(s), à l'exception des prestations où la suture est comprise, jusqu'au 18 ^e anniversaire	01-07-2022		N04	L21
375130	375141	no	yes	Honoraires complémentaires pour suture de plaine pendant une séance d'extraction(s) dentaire(s) ou ablation (section et extraction) de racine(s), à l'exception des prestations où la suture est comprise, par dent supplémentaire dans le même quadrant et durant la même séance, jusqu'au 18 ^e anniversaire	01-07-2022		N04	L12
375152	375163	no	yes	Honoraires complémentaires pour suture de plaine pendant une séance d'extraction(s) dentaire(s) ou ablation (section et extraction) de racine(s), à l'exception des prestations où la suture est comprise, par dent supplémentaire dans le même quadrant et durant la même séance, jusqu'au 18 ^e anniversaire	01-07-2022		N04	L8



This table shows the NIHDI nomenclature codes selected for this analysis, stating whether or not they were included in the analyses of services and expenditure, and giving, for each one, a description, dates of creation and deletion, where appropriate, their N group (in the NIHDI nomenclature) and their value.

B. Past history of nomenclature codes

Outpatient	Inpatient	Date	Label
301195	301206	01-01-2018	* Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, à partir du 53e anniversaire
301195	301206	01-07-2022	* Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, à partir du 50e anniversaire
301210	301221	01-01-2018	* Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, à partir du 18e anniversaire jusqu'au 53e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3 bis
301210	301221	01-07-2022	* Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, à partir du 18e anniversaire jusqu'au 50e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3bis
304850	304861	01-06-2010	* Extraction d'une dent à partir du 55e anniversaire
304850	304861	01-05-2014	* Extraction d'une dent à partir du 53e anniversaire
304850	304861	01-01-2018	* Extraction d'une dent, y compris la suture éventuelle, à partir du 53e anniversaire
304872	304883	01-06-2010	* Extraction d'une dent à partir du 55e anniversaire, par dent supplémentaire dans le même quadrant et au cours de la même séance
304872	304883	01-05-2014	* Extraction d'une dent à partir du 53e anniversaire, par dent supplémentaire dans le même quadrant et au cours de la même séance
304872	304883	01-01-2018	* Extraction d'une dent, y compris la suture éventuelle, à partir du 53e anniversaire, par dent supplémentaire dans le même quadrant et au cours de la même séance
304872	304883	01-07-2022	* Extraction d'une dent, à partir du 50e anniversaire, par dent supplémentaire dans le même quadrant et au cours de la même séance
304894	304905	01-06-2010	* Extraction d'une dent à partir du 18e anniversaire jusqu'au 55e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3 bis
304894	304905	01-05-2014	* Extraction d'une dent à partir du 18e anniversaire jusqu'au 53e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3 bis
304894	304905	01-01-2018	* Extraction d'une dent, y compris la suture éventuelle, à partir du 18e anniversaire jusqu'au 53e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3bis
304916	304920	01-06-2010	* Extraction d'une dent à partir du 18e anniversaire jusqu'au 55e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3 bis, par dent supplémentaire dans le même quadrant et au cours de la même séance
304916	304920	01-05-2014	* Extraction d'une dent à partir du 18e anniversaire jusqu'au 53e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3 bis, par dent supplémentaire dans le même quadrant et au cours de la même séance
304916	304920	01-01-2018	* Extraction d'une dent, y compris la suture éventuelle, à partir du 18e anniversaire jusqu'au 53e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3bis, par dent supplémentaire dans le même quadrant et au cours de la même séance
304916	304920	01-07-2022	* Extraction d'une dent, à partir du 18e anniversaire jusqu'au 50e anniversaire, dans le cas où le bénéficiaire répond à une des conditions de l'article 6, § 3bis, par dent supplémentaire dans le même quadrant et au cours de la même séance
371195	371206	01-01-2018	* Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, jusqu'au 18e anniversaire
371195	371206	01-07-2022	* Extraction chirurgicale d'une dent définitive, à l'exclusion des incisives, avec résection de l'os environnant et suture des lambeaux muqueux incisés, jusqu'au 18e anniversaire
374850	374861	01-05-2009	* Extraction d'une canine lactéale, d'une molaire lactéale ou d'une dent définitive, jusqu'au 18e anniversaire
374850	374861	01-01-2018	* Extraction d'une canine lactéale, d'une molaire lactéale ou d'une dent définitive, y compris la suture éventuelle, jusqu'au 18e anniversaire
374872	374883	01-05-2009	* Extraction d'une canine lactéale, d'une molaire lactéale ou d'une dent définitive jusqu'au 18e anniversaire, par dent supplémentaire dans le même quadrant et au cours de la même séance
374872	374883	01-01-2018	* Extraction d'une canine lactéale, d'une molaire lactéale ou d'une dent définitive, jusqu'au 18e anniversaire, par dent supplémentaire dans le même quadrant et au cours de la même séance
374872	374883	01-07-2022	* Extraction d'une canine lactéale, d'une molaire lactéale ou d'une dent définitive, jusqu'au 18e anniversaire, par dent supplémentaire dans le même quadrant et au cours de la même séance



This table displays the historic evolution of the definitions of the NIHDI-nomenclature codes taken into account for this analysis, if modifications were implemented during the period 2012-2022.

C. Source of data and analysis period

The data used in the analyses have been taken from the following databases:

Document N	for the utilisation rate and amount of expenses of insured persons (who meet the selection criteria) whose age, sex, preferential regime and residence are known 2012-2022
Document P	for the utilisation rate and amount of expenses of insured persons (who meet the selection criteria) by type of medical specialities in 2022
Document P, SHA, ADH	for the practice occurrences and analysis of patient care settings in 2021
-	-

Analysis period	2012-2022
-----------------	-----------



'N Documents' are monthly data sent by the sickness funds to NIHDI, within three months. These data show the number of services provided, dates and the fees involved. Every six months, these data are compiled by the insurers, which also add data on patients: age, gender, social category and district of residence. N Documents, however, cannot be used to analyse the combinations of services received by individual patients.

'P Documents' are six-monthly data sent by the sickness funds to NIHDI, within four months. These data show the services provided, the service-provider, the prescriber, the place of provision of service, and the hospital where patients were treated. P Documents can be used to monitor medical consumption and pricing, but not (yet) to analyse services per patient.

'Documents SHA, ADH' are sent annually and within six months by the insurer-organisations to the NIHDI. They include all the services provided respectively in day admission and standard hospitalisation, in general hospitals per hospital stay.

D. Specific selection criteria

Several filters may have been applied to the data, so that only one section of the population is considered in the analyses. If so, the filters used are shown in the table below:

FILTERS APPLIED TO DATA	
Sex	women and men
Age	all
-	-

E. Standardisation

The data are standardised before analysis per year, based on age, sex and preferential regime per arrondissement, province and region (standardization based on population in 2022).




Standardisation renders populations comparable in relation to one or several criteria. If a difference is observed between these populations, we can therefore assume that it is not due to the criteria covered by the standardisation process.

3. RESULTS

A. National standardised rate of use

	TOTAL
Average number of interventions per year	618.119
Standardised rate of use per 100 000 insured persons	5.338

B. Breakdown of nomenclature codes provided, by volume



See page 4 for details about the NIHDI nomenclature codes selected for analysis.

Note : The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis.

C. Specialisation of healthcare providers

Specialisation of the provider	Total providers	Concerned providers	% Providers	Median of H.C. services	Q3 of H.C. services	% Total H.C. services
Dentists	8024	7385	92%	38	69	63,76%
Stomatologists	647	628	97%	263	439	30,08%
Periodontologists	209	195	93%	75	150	3,48%
Dentists in training	1192	715	60%	13	31	2,57%
Other specialities	5357	32	1%	13	70	0,11%
Total	15429	8955	58%	39	75	100,00%



This table shows the following non-standardised data, by medical specialities (figures for the year 2022):

- The number of service-providers per specialisation who have recorded at least one service (the figures are exceptionally extrapolated from a single semester if an * is indicated in the header, otherwise the full year is taken into account);
- The number of service-providers recording services under the nomenclature codes selected for this analysis;
- The service-providers for these codes as a percentage of the total number of service-providers recording provision of at least one service;
- The median number and third quartile of services per service-provider (recording provision under these codes);
- The service percentage, i.e. the number of services recorded for this specialisation as a percentage of total services provided.

D. Specialisation of prescribers

Specialisation of the prescriber	Total prescribers	Concerned prescribers	% Prescribers	Median of prescriptions	Q3 of prescriptions	% Prescriptions
Not applicable	0	0	0%	0	0	100,00%



This table shows, in order, the following non-standardised data per specialities (figures for the year 2022):

- The number of prescribers who have prescribed at least one service (the figures are exceptionally extrapolated from a single semester if an * is indicated in the header, otherwise the full year is taken into account);
- The number of prescribers prescribing the nomenclature codes selected for this analysis;
- The prescribers prescribing these codes as a percentage of the number of prescribers prescribing at least one service;
- The median number and third quartile of services per prescriber (prescribing these codes);
- The percentage of services prescribed, i.e. the number of prescriptions issued for this specialisation as a percentage of total services prescribed.


E. Standardised rate of use by sex and age group

	TOTAL
<i>Average number of interventions per year</i>	618.119
Median age (years)	59
Mean age (years)	50,43
Max/Min Ratio of the median age (by district)	1,16
Percentage of women	50,61%

Max/Min Ratio:

The max/min ratio measures the dispersion of values. It is calculated as the ratio of the maximum value found for the variable, in all districts, to the minimum value. If this minimum value is equal to zero, the max/min ratio cannot be calculated, and is reported as 'NA' ('not applicable').

Dentists - Extraction (Adjusted)




Standardised rate of use per 100 000 insured persons, and coefficient of variation for the districts, by age group and sex, for the year 2022

This figure is made up of bar charts for each sex. The **coefficient of variation**, shown by the red line, measures the relative dispersion of the standardised rates of use observed for each district, by age group and sex (standard deviation divided by the mean). This line is shown in bold for age groups where the coefficient of variation can be validly interpreted (i.e. for age groups in which there are sufficient insured persons per district to allow for a proper comparison).


The left-hand vertical axis of the graph represents the standardised rate of use, and the right-hand axis the coefficient of variation. The horizontal axis shows the age groups. The horizontal dotted lines show the total values of the standardised rates of use (in blue) and of the coefficient of variation (in red).

Dentists - Extraction (Adjusted)



Comparison of the standardised rates of use by sex (per 100 000) in 2022

Dentists - Extraction (Adjusted)




Standardised rate of use per 100 000 insured persons, by sex and by province for the year 2022


This histogram shows standardised rates of use by province and by sex. The grey bars show the rates for men, while the green bars show the rates for women, for each province. The grey and green broken lines show the total standardised rates of use, again grey for men, green for women.

F. Standardised rate of use: hospital and outpatient care

	TOTAL
Average number of interventions per year	618.119
Percentage of out-patient care	98,87%
Max/min ratio of out-patient care percentage (by district)	1,02



This graph shows the percentage of outpatient services (including hospital day admissions), i.e. the number of outpatient services provided as a percentage of total services (outpatient and hospital stays). Besides the bar per region, there is a bar for the entire Belgian population. A dotted line also shows this overall ratio.




N.B.:

- The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis
- A complementary document to this chapter, about the handling of patients per health care sector, is [enclosed in this report](#) (cf. p.36)

G. Standardised rate of use by reimbursement scheme

	TOTAL
<i>Average number of interventions per year</i>	618.119
Percentage provided under the preferential reimbursement scheme	22,44%
Standardised rate of use with preferential reimbursement scheme (per 100 000)	5.826
Standardised rate of use without preferential reimbursement scheme (per 100 000)	5.276
Ratio Preferential scheme /General scheme	1,1

Dentists - Extraction (Adjusted)




This graph shows the standardised rates of use with (in red) and without (in grey) the preferential reimbursement scheme, by region and in total. The red and grey dotted lines show the overall standardised rates of use, with and without the preferential reimbursement scheme, respectively.

H. Trends in standardised rates of use

	TOTAL	Statistical significance
Average number of interventions per year	618.119	
Trend (2012-2022)	1,39%	*** (1,46%)
Trend (2012-2019)	1,08%	***
Trend (2019-2022)	2,12%	

These trends correspond to the average annual growth rate.

A non-significant statistical test indicates that the trend estimated by the model (in brackets) is stable, or that there is no break in the trend




This graph shows a coloured curve for each region and a black curve for the entire Belgian population. The x-axis shows the years, and the y-axis shows the standardised rate of use per 100 000 insured persons.

Note : The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis.

Trends in the standardised rate of use per 100 000 insured persons, by region

Dentists - Extraction (Adjusted)



This graph shows a colored line for each district and a black line for the entire Belgian population. The x-axis shows the years, and the y-axis shows the standardised rate of use per 100 000 insured persons.

To better highlight changes over time, the rates shown are **rolling averages** of the rates for the three years preceding the year in question (including the year itself).

The graph only shows the five districts with the highest average rates and the five districts with the lowest average rates over the last 3 years studied.

Note : The year 2020 was highlighted by a vertical dashed line, in order to draw the attention on the impact of the COVID-19 crisis.

	Rate of use	Annual increase			Structural break	
		2022 (per 10 ⁵ insured)	2012- 2022	2012- 2019	2019- 2022	
Provinces	West Flanders	5716,91	1,43%	1,36%	1,60%	NA
	East Flanders	5633,44	2,11%	1,95%	2,48%	NA
	Antwerp	5276,48	1,00%	1,00%	0,98%	NA
	Limburg	5208,37	0,94%	0,59%	1,76%	NA
	Flemish Brabant	5409,25	0,97%	0,60%	1,83%	NA
	Brussels	5357,51	1,21%	0,67%	2,49%	NA
	Walloon Brabant	5305,9	1,45%	1,01%	2,49%	NA
	Hainaut	4980,58	1,44%	1,16%	2,09%	NA
	Liège	5348,86	1,60%	0,77%	3,55%	NA
	Namur	5076,64	1,88%	1,28%	3,31%	NA
Regions	Luxembourg	4650,9	2,14%	1,26%	4,21%	NA
	Flanders	5451,6	1,31%	1,15%	1,68%	NA
	Brussels	5357,51	1,21%	0,67%	2,49%	NA
	Wallonia	5119,73	1,60%	1,04%	2,92%	NA
	TOTAL	5337,61	1,39%	1,08%	2,12%	***


Trends in the rates of use, by province and region

This table reports the standardised **rates of use** for the last year analysed (2022), as well as the average **rates of increase**, by province, by region and in total, for the entire period (2012-2022), for the last years (2019-2022) and for the period preceding the last years (2012-2019)

In order to find out whether the trend in the last years differs from that in the years before, a linear mixed model was fitted in two steps. In the first step a change in trend on the national level is tested. If this test is significant, in a second step, the model tests whether the difference in trend is significant for each province, region and at the national level. The data of 2020 are excluded from the models.

The significance of the test for a change in trend is reported in the Structural break column : * P-value ≤ 0.05 / ** P-value ≤ 0.01 / *** P-value ≤ 0.001 and NS for a non-significant result.

'NA' is shown where the nomenclature codes selected for the analysis have not been used for the entire last period or when the statistical tests cannot be evaluated.



Trend break assessment model by province – Regression lines

Regression lines per province showing a possibly different slope for the last years (2019-2022) compared to the years before (2012-2019).


Data of 2020 was excluded from this analysis, but is indicated on the graph for information.

I. Geographical variations in standardised rates of use

	TOTAL
<i>Average number of interventions per year</i>	618.119
Coefficient of Variation (2022)	7,64
Max/Min Ratio* of the standardised rates of use (by region)	1,06
Max/Min Ratio* of the standardised rates of use (by district)	1,39

Coefficient of Variation (2020-2022)	10,03
Coefficient of Variation (2012-2014)	9,34
<i>Statistically significant difference? (p ≤ 0.05)</i>	No

* An 'NA' result indicates a ratio which cannot be calculated, i.e. the minimum value = zero (cf. E. Standardised rate of use by sex and age group)



A **dot plot** is a distribution chart, which is useful for highlighting groups in the data, gaps in the distribution and outliers. Here, each dot represents the rate of use of a district, for its entire population or broken down by sex.

The rates are rounded to the nearest unit, ten, hundred, etc., depending on the value of the maximum rate, in order to better group the values.

The graph also shows a box with the 25th, 50th and 75th percentiles of the non-rounded standardised rates of use for all patients. The bottom line of the box represents the 25th percentile, while the upper line represents the 75th percentile. The line inside the box represents the 50th percentile.



Map showing distribution of standardised rates of use, by district

On this map of Belgium, thin lines show the boundaries of the districts, while thick lines show the provincial borders. The districts are coloured using a colour scale based on the level of rate of use in the district compared to the Belgian national rate (overall rate). This ratio is expressed as a percentage: e.g. 0% if the district rate is equal to the overall rate, 20% if the rate is 20% above the overall rate, and -20% if the rate is 20% below the overall rate. The percentages are calculated using the standardised rates of the last year analysed, and are displayed in bands of 20%. The following colour coding applies:

Colour	Category
Red	More than 50%
Orange	Between 30% and 50%
Light Orange	Between 10% and 30%
Yellow	Between -10% and 10%
Light Green	Between -30% and -10%
Dark Green	Between -50% and -30%
Very Dark Green	Less than -50%
White	Not used


N.B.: The interpretation of this map is to be done in parallel with [the graph in funnel plot \(p.28\)](#)



Map showing distribution of standardised expenditure, by district

On this map of Belgium, thin lines show the boundaries of the districts, while thick lines show the provincial borders. The districts are coloured using a colour scale based on the level of expenditure in the district compared to Belgian national (overall) expenditure. This ratio is expressed as a percentage: e.g. 0% if expenditure in the district is equal to the overall expenditure, 20% if it is 20% higher, and -20% if it is 20% lower. The percentages are calculated using the standardised expenditure of the last year analysed and are displayed in bands of 20%. The following colour coding applies:

Colour	Category
Red	More than 50%
Orange	Between 30% and 50%
Light Orange	Between 10% and 30%
Yellow	Between -10% and 10%
Light Green	Between -30% and -10%
Dark Green	Between -50% and -30%
Black	Less than -50%
White	No expenditure



**'Funnel plot' showing the standardised rates of use by district,
by the number of insured persons**

In this graph, the standardised rate of use in a district is positioned versus the size of its population. Besides the dots representing the districts, 95% and 99.7% **confidence intervals** are also shown on the graph. These are dependent of the size of the districts. The thicker horizontal line shows the national standardised rate of use. The outlier districts are identified as those districts that fall outside the 99.7% confidence intervals, the zone between the 95% and 99.7% confidence intervals being considered as "warning zone".

N.B.: The interpretation of this graph is to be done in parallel with the [map of the distribution of rates of use \(p.26\)](#)

J. Standardised healthcare expenditure borne by the insurance

	TOTAL
<i>Average number of interventions per year</i>	618.119
Average annual expenditure (€)	76.847.210
Average cost per intervention (€)	124,32
Average annual expenditure per insured (€)	6,64
Max/Min Ratio* of expenditure per insured (by region)	1,22
Max/Min Ratio* of expenditure per insured (by district)	1,94

* An 'NA' result indicates a ratio which cannot be calculated, i.e. the minimum value = zero (cf. E. Standardised rate of use by sex and age group)

		Standardised expenditure (per insured)
Provinces	West Flanders	7,41 €
Regions	East Flanders	7,29 €
Provinces	Antwerp	7,09 €
Regions	Limburg	7,16 €
Provinces	Flemish Brabant	6,82 €
Regions	Brussels	5,94 €
Provinces	Walloon Brabant	6,38 €
Regions	Hainaut	6,51 €
Provinces	Liège	4,93 €
Regions	Namur	6,35 €
Provinces	Luxembourg	4,74 €
Regions	Flanders	7,15 €
Regions	Brussels	5,94 €
Regions	Wallonia	5,89 €
	TOTAL	*** €

Regional and provincial distribution of standardised expenditure (2022)

Dentists - Extraction (Adjusted)

Nomenclature	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	Average annual growth rate
301151-301162	0,00	0,00	0,00	0,00	0,00	0,00	98,40	100,65	102,31	105,25	107,21	2,17%
301173-301184	0,00	0,00	0,00	0,00	0,00	0,00	100,86	104,65	105,94	106,26	107,30	1,56%
301195-301206	0,00	0,00	0,00	0,00	0,00	0,00	83,80	85,40	87,50	89,47	91,30	2,17%
301210-301221	0,00	0,00	0,00	0,00	0,00	0,00	86,38	88,60	90,85	92,52	94,09	2,16%
304850-304861	59,54	61,51	62,34	62,30	62,23	62,59	68,89	70,50	71,69	74,91	75,41	2,39%
304872-304883	42,77	44,31	44,87	44,88	44,56	44,46	47,03	48,26	49,43	50,20	51,95	1,96%
304894-304905	61,44	62,98	64,31	64,68	64,39	64,39	69,92	72,55	72,66	76,03	76,38	2,20%
304916-304920	45,93	47,07	48,61	48,91	47,93	47,62	50,31	52,38	52,66	53,16	54,96	1,81%
304975-304986	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	75,03	NA
304990-305001	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	75,41	NA
312410-312421	380,75	383,28	387,87	386,78	386,94	390,04	395,08	396,06	400,65	402,92	408,32	0,70%
312432-312443	353,90	357,73	361,93	362,49	364,43	366,73	373,05	373,56	375,89	380,29	387,65	0,91%
317214-317225	176,18	181,12	181,53	183,85	185,22	185,01	185,76	190,51	192,32	196,89	190,29	0,77%
317236-317240	184,70	188,08	191,91	194,50	193,20	195,16	195,93	195,84	199,13	198,27	197,70	0,68%
317251-317262	91,26	91,94	93,30	93,31	93,86	94,09	95,80	96,01	96,16	97,96	98,31	0,75%
317376-317380	229,70	228,45	213,76	239,99	242,28	235,48	237,70	220,23	260,61	183,69	190,78	-1,84%
317391-317402	111,71	107,42	111,77	104,21	101,58	107,05	99,62	104,30	100,71	108,71	102,01	-0,90%
371151-371162	0,00	0,00	0,00	0,00	0,00	0,00	115,19	116,67	120,29	122,41	122,63	1,58%
371195-371206	0,00	0,00	0,00	0,00	0,00	0,00	129,56	131,28	133,61	135,68	137,39	1,48%
374850-374861	73,30	75,67	76,85	76,91	76,92	77,74	83,97	85,65	86,50	88,90	88,90	1,95%
374872-374883	51,96	53,66	54,54	54,51	54,52	55,52	58,16	59,44	60,21	60,75	62,18	1,81%
374975-374986	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	0,00	89,07	NA

Change over time in expenditure, by service and by nomenclature code

4. KEY DATA SUMMARY

		TOTAL
PROVIDERS & PRESCRIBERS		
Main healthcare providers:	Dentists	63,76%
Main prescribers:	Not applicable	-
RATE OF USE		
Number of interventions (per year)		618.119
Standardised rate of use (per 100 000 insured persons)		5337,61
≥ 2 occurrences per patient (2021) ⁴		45,9%
Percentage of outpatient care		98,87%
POPULATION		
Median age		59 years
Max/min ratio ⁵ of the median age (by district)		1,16
Percentage of women		50,61%
Ratio Preferential rate/General rate		1,1
TRENDS		
Trend ⁶ (2012-2022)	1,39%	***
Trend ⁶ (2012-2019)	1,08%	***
Trend ⁶ (2019-2022)	2,12%	
GEOGRAPHICAL VARIATIONS		
Coefficient of variation ⁶ (2012-2014)	9,34	
Coefficient of variation ⁶ (2020-2022)	10,03	NS
Max/min ⁵ Ratio of number of interventions ⁶ (per 100 000 insured persons, by region)	1,06	
Max/min Ratio ⁵ of number of interventions ⁶ (per 100 000 insured persons, by district)	1,39	
DIRECT EXPENDITURE		
Average annual expenditure	76.847.210 €	
Average annual expenditure per insured	6,64 €	
Max/Min Ratio ⁵ of expenditure per insured (by region)	1,22	
Max/Min Ratio ⁵ of expenditure per insured (by district)	1,94	
Average cost of interventions	124,32 €	
CODING VARIATIONS & PRACTICE ALTERNATIVES⁴		
Variations in practice coding ⁶ (by province)	Yes	***
Variations in the choice of practice alternatives ⁶ (by province)	Yes	***

⁴ More detailed results are shown in a document enclosed to this report.

⁵ An 'NA' result indicates a ratio, which cannot be calculated, i.e. the minimum value equals zero.

⁶ If the result(s) show(s) a significant difference, the level of statistical significance is symbolized by one to three asterisks (increasingly significant). Otherwise, NS is displayed (not significant). 'NA' indicates the test is not applicable.

5. APPENDICES

A. Analysis of variance (ANOVA), except Brussels

Statistical significance of the differences observed in 2022		
<i>By region?</i>	Yes	***
<i>By sex?</i>	Yes	***
<i>By reimbursement scheme?</i>	Yes	***
<i>By sex and per region?</i>	Yes	***
<i>By reimbursement scheme and per region?</i>	No	NS
<i>By sex and per reimbursement scheme?</i>	No	NS
<i>By sex and reimbursement scheme and per region?</i>	Yes	*


In order to be able to assess the significance of the observed differences, a linear mixed ANOVA model was fitted to the data of all districts of the Walloon and Flemish regions, after standardising for age. The model has region, sex and reimbursement scheme as fixed effects and also contains all two-way and three-way interactions between these effects.

In order to interpret the model correctly, first the three-way interaction should be evaluated, followed by the two-way interactions and finally by the main effects. If the three-way interaction is significant, the interpretation of the model should be done at this level only and the two-way interactions and main effects should not be interpreted. If the three-way interaction is not significant, the two-way interactions are evaluated. Every main effects that appears in a significant interaction should be interpreted at the level of the interaction and not at the level of that main effect. Main effects can only be interpreted directly if they don't appear in a significant interaction.

The **asterisks** represent the level of statistical significance of the tests: * P-value ≤ 0,05 / ** P-value ≤ 0,01 / *** P-value ≤ 0,001 or NS for a non-significant result.

B. Frequency of practice occurrences

Frequency	Per year	Per day
2 occurrences	19,2%	19,1%
≥ 3 occurrences	26,7%	22,1%
≥ 2 occurrences	45,9%	41,2%



Distribution of practice recurrences per year (2021)

Some practices may be billed several times for the same patient in the same year or even on the same day. This may be due to a **repetition of the practice**, but also to an anatomical effect, which may lead, depending on the organ concerned, to performing the same practice **bilaterally**, which may therefore cause a double occurrence on the same day.




In order to interpret the results per day validly, it is useful to note that the same patient may be counted several times if, for example, he or she has received two identical services simultaneously, twice a year.

These frequency analyses of occurrences are carried out over the year **2021** using the following databases: Documents P, ADH and SHA.


Values « **n.a.** » are indicated if the data were not available at the time of this report.

Dentists - Extraction (Adjusted)



Frequency of practice occurrences by province and variation vs national value (2021)

C. Patient care settings



█ Inpatient █ One-day
█ Outpatient (clinic) █ Outpatient (private)


Distribution of patient care settings in 2021

Care Settings	
Outpatient (private)	70,6%
Outpatient (polyclinic)	7,6%
(Day) Hospital	21,1%
Hospital (stay)	0,6%

In addition to the chapter on [standardised inpatient and outpatient use rates](#) (see p.16), the analysis of patient care settings can be refined by identifying the outpatient (private and polyclinic) and inpatient (day or standard hospitalisation) sub-sectors.

These analyses are carried out over the year **2021** using the following databases: Documents P, ADH and SHA.


Values « n.a. » are indicated if the data were not available at the time of this report.



Distribution of patient care settings by province (2021)

D. Coding variations and practice alternatives

➔ Variations in coding:




[Due to the large number of nomenclature codes selected for this analysis, we cannot include them in the legend here. We therefore invite you to read the details concerning them on page 4 of this report.]

Significance	By region	By province
Use of Nomenclature codes ⁷	***	***

⁷ The calculation of significance is carried out here by comparing the geographical differences in the use of the different nomenclature codes to code the practice.

The asterisks represent the level of statistical significance of Chi-square test: * P-value ≤ 0,05 / ** P-value ≤ 0,01 / *** P-value ≤ 0,001. NS and NA respectively indicate that the variations are not significant or not applicable.

➔ Variations in practice alternatives:



Breakdown of choice for practice alternatives


Significance	By region	By province
Choice of Practice alternatives	***	***

Combined codes	Groupings
301151-301162	1_non_surgical
301173-301184	1_non_surgical
301195-301206	2_surgical
301210-301221	2_surgical
304850-304861	1_non_surgical
304872-304883	1_non_surgical
304894-304905	1_non_surgical
304916-304920	1_non_surgical
304975-304986	1_non_surgical
304990-305001	1_non_surgical
312410-312421	3_Embbeded_Tooth
312432-312443	3_Embbeded_Tooth
317214-317225	4_hospital
317236-317240	3_Embbeded_Tooth
317251-317262	4_hospital
317376-317380	4_hospital
317391-317402	4_hospital
371151-371162	1_non_surgical
371195-371206	2_surgical
374850-374861	1_non_surgical
374872-374883	1_non_surgical
374975-374986	1_non_surgical



According to the nature of the practice and the analytical tools available, it may be possible to identify and define alternatives for carrying out the practice. In this case, the nomenclature codes defined for the analysis of the practice are grouped together with the aim of analysing whether or not the choices of these alternatives are homogeneous across the territory. The calculation of significance displayed in the table is carried out by comparing these groups of codes with each other.

→ Variation in the use of Global Payment with Standardisation (GPS) :



Pseudocodes	Label
-------------	-------

Breakdown by volume of pseudocodes of GPS

Significance	By region	By province
Use of Global Payment with Standardisation ⁸	NS	NS

⁸ The calculation of significance is carried out here by comparing the use of Global Payment with Standardisation as a whole compared to the non-use of these packages.